



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

59. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Show that the tangent plane at any point of the surface $a^2x^2 + b^2y^2 + c^2z^2 = 2bcyz + 2acxz + 2abxy$ intersects the surface $ayz + bzx + cxy = 0$ in two straight lines at right angles to one another.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

47. Proposed by Prof. J. SCHEFFER, A. M., Hagerstown, Maryland.


The floor of a vault forms a square, and all sections parallel to it are squares. The two vertical sections through the middle points of the opposite sides of the floor are equal semi-circles. Find the convex surface and the volume of the vault.

I. Solution by C. W. M. BLACK, Professor of Mathematics in Wesleyan Academy, Wilbraham, Massachusetts; O. W. ANTHONY, M. Sc., Professor of Mathematics in New Windsor College, New Windsor, Maryland; A. H. HOLMES, Brunswick, Maine, and the PROPOSER.

Let $ABCD$ represent the base square, side $= 2a$, and KEI and GFH the two equal semi-circles, radius $= a$. Let $LMNO$ be another square parallel to the base square, and at the distance $PE = x$ from it. The area of $LMNO$ is $= 4(a^2 - x^2)$,

$$\therefore \text{Vol.} = 4 \int_0^a (a^2 - x^2) dx = \frac{8}{3} a^3.$$

Denoting $\angle PEQ$ by θ , we have for the surface



$$\int_0^{\frac{1}{2}\pi} 8a \cos \theta d(\theta) = 8a^2. \quad \text{Or for the volume, } dV = 4a^2 \cos^2 \theta dx, \text{ where } x \text{ is}$$

the vertical distance, $x = a \sin \theta$; $dx = a \cos \theta d\theta$.

$$\therefore V = 4a^3 \int_0^{\frac{1}{2}\pi} \cos^3 \theta d\theta = a^3 \int_0^{\frac{1}{2}\pi} (\cos 3\theta + 3\cos \theta) d\theta = \frac{8a^3}{3}.$$

II. Solution by BENJ. F. YANNEY, A. M., Professor of Mathematics in Mount Union College, Alliance, Ohio.

The convex surface of the vault is equivalent to the surface of a right cir-

cular cylinder intercepted by another right circular cylinder, their axes intersecting at right angles, the two cylinders being equal, and the diameter of each equal to that of the vertical sections of the vault.

\therefore Letting the radius $= a$, $S = 8a \int_0^a \int_0^{\sqrt{a^2-x^2}} \frac{xdy}{\sqrt{a^2-x^2}} = 8a^2$, the equa-

tions of the cylinders being $x^2 + z^2 = a^2$, and $x^2 + y^2 = a^2$.

The volume is equivalent to that of four wedges cut from the cylinder, $x^2 + y^2 = a^2$, by the planes, $z=0$, and $z=x$.

$$\therefore V = 8 \int_0^a \int_0^{\sqrt{a^2-x^2}} \int_0^x dx dy dz = \frac{8a^3}{3}.$$

Also solved by E. L. SHERWOOD and G. B. M. ZERR.

48. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.


I have a circular section basin 12 inches in perpendicular height; the diameters are as follows: At base, 2 inches; one inch perpendicular height, 6 inches; two inches perpendicular height, 18 inches; three inches perpendicular height, 54 inches; and so on, the diameter being trebled for every inch in height. After a rain the water in the basin is six inches deep, what was the rainfall?

I. Solution by E. L. SHERWOOD, A. M., Professor of Mathematics, Mississippi Normal College, Houston, Mississippi; C. W. M. BLACK, A. M., Professor of Mathematics in Wesleyan Academy, Wilbraham, Massachusetts, and the PROPOSER.

The basin is generated by revolving the curve $x=3^y$ about the axis of y .

$$\therefore \text{Volume of water} = \pi \int_0^6 x^2 dy = \pi \int_0^6 3^{2y} dy.$$

$$\therefore V = \pi \frac{3^{12} - 1}{2 \log 3} = \frac{531440\pi}{2 \log 3}.$$



Let x = depth of rain-fall, then since radius of top of basin $= 3^{12}$, $V = \pi 3^{24} x$.

$$\therefore x = \frac{565720}{282429536481 \log 3} = .00000086 \text{ inches.}$$

II. Solution by O. W. ANTHONY, M. Sc., Professor of Mathematics in New Windsor College, New Windsor, Maryland.

Call x the length of any radius, and y the vertical distance, y being 1 at the bottom of the basin. Then the equation of side of basin is $x=3^{y-1}$,

$$dV = \pi x^2 dy, \quad V = \pi \int_1^y 3^{2y-2} dy = \frac{\pi [3^{12} - 1]}{3 \log 3}.$$

The radius of upper base $= 3^{12}$. Call R the rainfall, then